



U.S. EPA  
OSWER #9285.7-37  
March 2003

**TRW RECOMMENDATIONS FOR  
PERFORMING HUMAN HEALTH RISK  
ANALYSIS ON SMALL ARMS SHOOTING  
RANGES**

Office of Solid Waste and Emergency Response  
U.S. Environmental Protection Agency  
Washington, DC 20460

## **NOTICE**

This document provides guidance to EPA staff. It also provides guidance to the public and to the regulated community on how EPA intends to exercise its discretion in implementing the National Contingency Plan. The guidance is designed to implement national policy on these issues. The document does not, however, substitute for EPA's statutes or regulations, nor is it a regulation itself. Thus, it cannot impose legally-binding requirements on EPA, States, or the regulated community, and may not apply to a particular situation based upon the circumstances. EPA may change this guidance in the future, as appropriate.

**U.S. ENVIRONMENTAL PROTECTION AGENCY  
TECHNICAL REVIEW WORKGROUP FOR LEAD**

The Technical Review Workgroup for Lead (TRW) is an interoffice workgroup convened by the U.S. EPA Office of Solid Waste and Emergency Response/Office of Emergency and Remedial Response (OSWER/OERR).

**CO-CHAIRPERSONS**

**Region 8**

Jim Luey  
Denver, CO

**NCEA/Washington**

Paul White

**MEMBERS**

**Region 1**

Mary Ballew  
Boston, MA

**Region 10**

Marc Stifelman  
Seattle, WA

**Region 2**

Mark Maddaloni  
New York, NY

**NCEA/Washington**

Karen Hogan

**Region 3**

Linda Watson  
Philadelphia, PA

**NCEA/Cincinnati**

Harlal Choudhury

**Region 4**

Kevin Koporec  
Atlanta, GA

**NCEA/Research Triangle Park**

Robert Elias

**Region 5**

Patricia VanLeeuwen  
Chicago, IL

**OERR Mentor**

Larry Zaragoza  
Office of Emergency and Remedial Response  
Washington, DC

**Region 6**

Ghassan Khoury  
Dallas, TX

**Executive Secretary**

Richard Troast  
Office of Emergency and Remedial Response  
Washington, DC

**Region 7**

Michael Beringer  
Kansas City, KS

**Associate**

Scott Everett  
Department of Environmental Quality  
Salt Lake City, UT



## **EXECUTIVE SUMMARY**

Past management practices at small arms outdoor shooting ranges (subsequently referred to as ‘ranges’) allowed the spent ammunition to accumulate on site. Many range operators now recognize the risk posed to humans and the environment by spent lead ammunition and have implemented programs to manage and recycle lead shot and bullets. In accordance with its mission to provide scientifically sound and consistent guidance on lead risk assessment, the TRW has prepared this document to provide guidance and recommendations for performing risk assessment on land currently or formerly used as ranges. This document supplements Region 2’s [Best Management Practices for Lead at Outdoor Shooting Ranges](#) (U.S. EPA, 2001a), which serves as national guidance on the management of lead at ranges to minimize the environmental impact of spent lead ammunition (U.S. EPA, 2001b).

As used in this document, the term ‘small arms’ includes rifles, handguns (pistols), shotguns, submachine guns, and machine guns (NRA, 1999, p. I-8). Ranges can be divided into high velocity shooting ranges, where target shooting with pistols and rifles occurs, and low velocity shooting ranges where shotguns are used (i.e., skeet, trap, and sporting clay ranges). Lead bullets and fragments at pistol and rifle ranges are typically contained in a relatively small, well-defined area, or volume, of sand and/or soil. The shotfall zones at skeet and trap ranges may cover 10–50 acres or more, depending upon the layout of the range.

This document contains brief discussions of the regulatory background for outdoor shooting ranges and the toxicology of lead on humans, an operational and physical description of the different types of outdoor shooting ranges, and the fate of spent lead ammunition in the environment and its bioavailability. This document provides recommendations on how the Integrated Exposure Uptake Biokinetic model and the Adult Lead model can be used to predict the risk to human health from spent lead ammunition on small arms shooting ranges.

## **BACKGROUND AND PURPOSE**

There are approximately 9,000 outdoor small arms shooting ranges (subsequently referred to as ‘ranges’) in the United States, excluding those located on military sites (U.S. EPA, 2001a). Millions of pounds of lead are discharged annually at these ranges (U.S. EPA, 2001a). In the past, the common practice at ranges was to allow the spent ammunition to accumulate on site. Many range operators now recognize the risk posed to humans and the environment by the lead in spent ammunition and have implemented programs to manage and recycle lead shot and bullets.

Given the large number of ranges in the United States and their potential risk to humans, the Technical Review Workgroup for Lead (TRW) has prepared this document to assist Regional and State risk assessors and project managers in performing risk assessments at range sites. The proper management of lead at shooting ranges is addressed in an EPA Region 2 document entitled “[Best Management Practices for Lead at Outdoor Shooting Ranges](#)” (U.S. EPA, 2001a). This white paper contains the TRW’s recommendations on how to collect data that will be used to provide site specific information when using the Integrated Exposure Uptake Biokinetic ([IEUBK](#)) Model and the Adult Lead Model ([ALM](#)) (U.S. EPA, 2002c). The focus of this document is on formerly used ranges, although exposure to active ranges may occur and is also discussed.

## **INTRODUCTION**

As used in this document, the term ‘small arms’ includes rifles, handguns (pistols), shotguns, submachine guns, and machine guns (NRA, 1999, p. I-8). Spent ammunition on ranges is not regulated as solid/hazardous waste unless it is discarded (abandoned) and left to accumulate for a long period of time (U.S. EPA, 2001a). Furthermore, it is not regulated if the spent ammunition is recovered or reclaimed on a regular basis (U.S. EPA, 2001a). However, if the range poses an imminent or substantial danger to health or the environment it can be addressed through Resource Conservation and Recovery Act (RCRA) (U.S. EPA, 2001a).

The intake of lead has a wide variety of effects on humans. Adults and children exposed to lead are susceptible to neurotoxic effects (ATSDR, 1999). Lead may also increase blood pressure and cause anemia; high levels of exposure to lead may damage the brain and kidneys and even cause death (ATSDR, 1999; U.S. EPA, 2002b). High levels of exposure may cause miscarriages, retard fetal development, and damage the organs responsible for sperm production (ATSDR, 1999; U.S. EPA, 2002b). Other effects of lead exposure include irritability, poor muscle coordination, muscle and joint pain, memory and concentration problems, digestive problems, and hearing and vision impairment (U.S. EPA, 2001a, 2002b). In children, lead exposure can cause behavioral and learning problems, hearing problems, impairment of vision and motor skills, hyperactivity, and developmental delays (ATSDR, 1999; U.S. EPA, 2001a, 2002b). Blood lead concentrations of 10 µg/dL or less have been associated with adverse health effects in children (U.S. EPA, 1986a; CDC, 1991). It is EPA policy to limit exposure to lead such that the probability of a typical (or hypothetical) child, or group of similarly exposed children, having or exceeding the 10 µg/dL blood lead concentration is less than 5% (U.S. EPA, 1994a).

Other chemicals of potential concern at shooting ranges include arsenic and antimony (components of

ammunition), nickel (coating on some lead shot), copper, zinc, strontium, and magnesium (present in tracer munitions that are used in machine guns), and polycyclic aromatic hydrocarbons (present in petroleum 'pitch' found in clay targets used at skeet and trap ranges and in 'wadding' from shotgun shells) (Jorgensen and Willems, 1987; EEA, 1992; EA, 1995; Peddicord and LaKind, 2000). Lead shot contains primarily lead (97%), antimony (2%), arsenic (0.5%), and sometimes nickel (0.5%) (Jorgensen and Willems, 1987; Lin et al., 1995). The crust material surrounding lead shot contains between 0.5 and 2.0% antimony, 0.15% nickel, and trace amounts of arsenic (Jorgensen and Willems, 1987). Lead bullets are composed of 90–99% lead, 1–10.5% antimony, and 0.1% copper (EA, 1996). Baer et al. (1995) determined that clay targets ('pigeons') contain approximately 2/3 dolomitic sandstone and 1/3 pitch; painted targets also contain approximately 1% fluorescent paint.

Due to a ban on the use of lead shot for waterfowl hunting and in waterfowl production areas by the U.S. Fish and Wildlife Service (U.S. FWS, 1999, 2001), other less toxic materials have been introduced, such as bismuth, steel, tungsten/iron, and tungsten polymers (a list is provided in EPA, 2001a, Appendix B). However, due to the higher cost of shot that is manufactured with the less toxic material, lead shot continues to be the most commonly used material on skeet and trap shooting ranges (U.S. EPA, 2001b).

#### **TYPES OF SHOOTING RANGES**

Ranges can be divided into high velocity shooting ranges, where target shooting with pistols and rifles occurs, and low velocity shooting ranges where shotguns are used (i.e., skeet, trap, and sporting clay ranges). Appendix A contains a list of on-line sources of additional information for small arms outdoor shooting ranges.

#### **High Velocity (Pistol and Rifle) Shooting Ranges**

High velocity ranges consist of a firing line, targets, backstop (to contain bullets and fragments), side berms (to contain ricochets), and ground and overhead baffles (to contain 'short' and 'long' shots, respectively) (Vargas, 1996). Not all ranges, particularly older ranges, will include all these components. Typical lengths for ranges vary between 25 yards and 200 meters (642 yards) (Vargas, 1996); widths depend on the number of firing stations.

The backstop is required to contain the bullets and bullet fragments. Backstops traditionally consist of earthen berms, typically between 15 and 25 feet high. More recently, earthen berms have been replaced with sand traps, steel traps, and rubber traps (U.S. EPA, 2001a). Other innovations have been developed for backstops, such as Shock Absorbing Concrete (SACON) that has been used on some Department of Defense (DOD) ranges since the 1980s (U.S. EPA, 2001a). The purpose of the newer types of backstops/bullet containment devices is to facilitate the collection of bullets and bullet fragments, which substantially reduces the amount of contaminated media generated by the range.

#### **Low Velocity (Shotgun) Shooting Ranges**

Shotguns are used to shoot clay targets ('pigeons' or 'birds') on skeet, trap, and sporting clay ranges. In many cases, skeet and trap shooting takes place in one range. A typical trap range consists of five-shooting positions and one structure, the 'traphouse', from which the targets are thrown by a machine

called a 'trap'. The angle at which the targets are thrown varies within an arc of 45 degrees (in a horizontal plane). The shooting positions are located 16 yards from the traphouse (Capital Trap Club, 2001a). At skeet ranges, targets are released from two structures, the 'high house' and the 'low house.' There are eight-shooting positions arranged along an arc between the two houses. At the top of the arc, the shooter is approximately 30 yards from the line that connects the two houses (Capital Trap Club, 2001b). The actual layout of skeet and trap ranges varies widely between sites (e.g., EA, 1995; E&E, 1997; Murray et al., 1997).

The size and shape of the shotfall zone is a function of the layout of the site, and ranges from rectangular for sites with multiple ranges located next to each other (e.g., E&E, 1997), to semi-circular for sites with one range (e.g., EA, 1995). The outfall zone from trap shooting will tend to be less than for skeet shooting due to the angle at which shooting occurs. In skeet shooting, the targets are thrown overhead and the shooting angle is approximately 45 degrees from the horizontal. Targets are released much closer to the ground in trap shooting; the shooting angle is approximately horizontal. Another factor that affects the distance the shot will travel is the size of the shot used. When the shooting angle is approximately horizontal, the maximum distance shot will travel varies from 198 yards for No. 8 shot to 330 yards for No. 2 shot (Baldwin, 1994). Number 6 shot will cover an area between 300 and 700 feet from the shooting position when the shooting angle is level; if released from an angle of 40 degrees from the horizontal, the shot will drop between 400 and 900 feet from the shooting position (Baldwin, 1994).

#### **FATE OF LEAD AMMUNITION IN THE ENVIRONMENT AND BIOAVAILABILITY**

Lead ammunition oxidizes in the environment, forming a crust around the shot; this crust contains lead carbonates and sulfates (Jorgensen and Willems, 1987; Manninen and Tanskanen, 1993; Lin et al., 1995; EA, 1996; Murray et al., 1997). The predominate lead carbonates that have been found in the crust material include hydrocerussite ( $Pb_3(CO_3)_2(OH)_2$ ) and cerussite ( $PbCO_3$ ); the predominate lead sulfate compound is anglesite ( $PbSO_4$ ). The rate of oxidation depends upon several environmental factors including: oxidation/reduction potential, ionic strength, pH, oxygen content of the soil and the presence of compounds (e.g., phosphate) that may inhibit oxidation (Jorgensen and Willems, 1987; EA, 1996). At some point, the presence of the crust material appears to inhibit the further weathering of the ammunition (Jorgensen and Willems, 1987). While metallic lead is insoluble under typical environmental conditions, lead is released to the environment through the dissolution of the lead compounds found in the crust material (Jorgensen and Willems, 1987; Manninen and Tanskanen, 1993; Lin et al., 1995; EA, 1996; Murray et al., 1997). The solubility of the lead compounds is affected by pH, eH, the presence of carbonate, sulfate, sulfide, phosphate and chloride, and the organic matter content of the soil (Jorgensen and Willems, 1987; Manninen and Tanskanen, 1993; Lin et al., 1995; EA, 1996; Murray, 1997).

Site-specific environmental factors that affect weathering rates of lead on shooting ranges include the amount of precipitation, pH of rain water, slope of the ground surface, amount of organic material present on the ground surface (e.g., leaves, peat) and soil type (U.S. EPA, 2001a). In general, weathering of lead ammunition will increase with increasing precipitation amounts, increase with decreasing pH, increase with increasing chloride concentration, decrease with increasing ground slope



(due to decrease in contact time between precipitation and ammunition) and increase with increasing organic matter cover (U.S. EPA, 2001a). Disturbance of the soil (e.g., soil cultivation of agricultural fields) may increase the decomposition of lead ammunition (Jorgensen and Willems, 1987). Bundy et al. (1996) found high corrosion rates were negatively correlated with corrosion potential and soil resistance, two characteristics of the soil environment that can be measured in the field.

The weathering rate of small arms ammunition may be affected by the presence of a copper 'jacket' or metal casing that surrounds the lead core of some ammunition (Major, 2003). The contact between the different types of metals may create a galvanic couple that increases the rate of corrosion of the ammunition if the moisture content of the surrounding soil is sufficiently high. However, on some sites, jacketed bullets do not appear to corrode at a faster rate than unjacketed bullets (Hall, 2002). Jacketed bullets are common on small arms ranges that are located on DOD sites because the military is required to use jacketed bullets; jacketed bullets are much less common on non-military ranges (Hall, 2002). Bullet jackets typically contain (by weight) 89–95% copper, 0.03–0.05 % lead, 0.05% iron, and 5–10 % zinc (Battelle, 1997).

The bioavailability of lead compounds varies greatly from lead sulfates, which have relatively low bioavailability (<25% bioavailable), to lead carbonates (>75% bioavailable) (Henningsen et al., 1998). The overall bioavailability of lead at shooting ranges depends upon the relative amounts of lead carbonates and lead sulfates that are present in the soil. Equilibrium diagrams (i.e., eH-pH diagrams) predict lead sulfates to be the dominant form of lead at pHs <5.3, carbonates to be the dominant form at pHs between 5.3 and 8.5, and lead hydroxides to dominate at pHs >8.5 (EA, 1996). The speciation found at a particular site will also vary depending upon the amount of carbonate and sulfate present in the soil (EA, 1996).

#### **SITE CHARACTERIZATION AND RISK ASSESSMENT**

An important difference between high velocity ranges and low velocity ranges, with respect to risk assessment, is the size of the areas impacted by each type of range. Lead bullets and fragments at pistol and rifle ranges are typically contained in a relatively small, well-defined area, or volume, of sand and/or soil. Very little contamination may be found at a well-designed range equipped with bullet traps, although some traps and targets (e.g., steel targets) may generate lead dust and particulates that can be transported by air and water and contaminate the surrounding area.

Due to the relatively small size of typical pistol and rifle ranges, the risks posed to human and ecological receptors are typically low. Exceptions to this would include former ranges that are planned for development, or are currently used for activities that could result in exposure to human and ecological receptors. Risks to human and ecological receptors from exposure to lead and other contaminants at skeet and trap ranges may be moderate to severe, due to the size of the shot outfall area. In some cases, the outfall areas are located within or near wetlands and surface water, which tends to increase the risks to ecological receptors, particularly waterfowl (EA, 1995; E&E, 1997; U.S. EPA, 2001a).

The difference between the types of exposures at high velocity and low velocity ranges also warrant consideration in the exposure assessment. These differences are due to the nature of the lead contamination in soil. At high velocity ranges, in addition to whole bullets and fragments, small particles of lead will be present due to the partial disintegration of the bullets upon impact with the targets and soil and rock particles in the berms (EA, 1996). Less disintegration of the bullets can be expected where fully jacketed and partially jacketed bullets are used (e.g., military ranges). At low velocity ranges, the lead shot will tend to be less fragmented due to the lower velocities. Although lead shot does break down in the environment, complete decomposition of the lead is a slow process that may take 30–300 years, depending upon site conditions (Jorgensen and Willems, 1987; EA, 1996).

### **Exposure Scenarios and Pathways**

Land use adjacent to and near shooting ranges should be considered when developing current and future exposure scenarios for a site. Under the current land use scenario, the potentially exposed human populations of particular concern at an operating range are residents of adjacent residential properties, residents and farm workers on adjacent agricultural properties, and workers who are employed on adjacent commercial properties. Other receptors include trespassers who use the site for recreational purposes such as fishing, hunting, and hiking (Peddicord and LaKind, 2000; U.S. EPA, 2001a), as well as other recreational users when the range is located on or within an area that is used for recreational activities other than target shooting (e.g., multi-use parks). Under future land use scenarios, the potentially exposed population depends upon the intended or actual land use, which may include residential, agricultural, commercial, or industrial uses.

The potentially exposed populations under both future and current land use scenarios that should be considered are:

- Residential land use:
  - children under the age of 7 years old
  - adults
- Agricultural land use (farm family or subsistence farm family):
  - children under the age of 7 years old
  - adults
  - farm workers
- Commercial and industrial land use:
  - adults
  - trespassers
  - maintenance staff/construction workers who may be exposed during invasive work (e.g., excavating trenches to install/repair utilities)

The main pathway for human exposure to lead at shooting ranges is through incidental ingestion of contaminated soil (Peddicord and LaKind, 2000; U.S. EPA, 2001a). The ingestion of site-raised meat (beef, pork, chicken) and fruits and vegetables contaminated with lead dust may also pose a risk for local residents and especially for local farm families; these pathways must be evaluated in the latter scenario. Grazing farm animals may ingest and bioaccumulate large quantities of lead [Braun et al., 1997]. In addition, the inhalation of dust/soil particles may be a potential pathway depending upon

site conditions, particularly during activities that involve the excavation of soil (e.g., during maintenance and construction work), or during agricultural activities (e.g., tilling, planting, and plowing) that may release clouds of dust. Hunters (and poachers) may be exposed via consumption of lead-contaminated wildlife (EA, 1995; E&E, 1997; Peddicord and LaKind, 2000; U.S. EPA, 2001a). Children who exhibit pica behavior may be at risk from exposure to formerly used shooting range sites; however, the pica child will not be considered in this document. In most cases, the primary receptor of concern will be children and farmers who are potentially exposed to formerly used ranges and adjacent contaminated areas. (The subsistence farm family ingestion rate of relevant meat or site-raised crops is assumed to comprise a large part of the diet; data is currently available in the Exposure Factors Handbook to evaluate this scenario and the EPA Default Exposure Factors Workgroup is developing default values as well.)

Ecological receptors of concern at shooting ranges include invertebrates, fish, mammals and birds, particularly waterfowl. Pathways include the incidental ingestion of soil, intentional ingestion of lead fragments as grit, and the ingestion of contaminated food items. Risk assessments at shooting ranges have predicted unacceptable levels of risk from lead for raptors, and small (e.g., mice) and large (e.g., fox and deer) terrestrial mammals (EA, 1995; Peddicord and LaKind, 2000). The highest risks from lead have been predicted for small mammals and birds that ingest lead shot incidentally while feeding, or intentionally as grit (EA, 1995; Peddicord and LaKind, 2000). The predicted adverse effects of lead on small mammals and birds, are supported by the literature, which has shown mortality may result from the ingestion of one lead pellet (e.g., Ma, 1989; Roscoe et al., 1989; Hoffman et al., 2000; Vyas et al., 2000). Additional information is available from the EPA [ECOTOX](#) database (U.S. EPA, 2002a). Consultation with an ecotoxicologist is recommended when planning an ecological risk assessment.

### **Soil Sampling Strategies and Recommendations**

This section is divided into three subsections: 1) General Sampling Strategy, 2) Sample Collection at High Velocity Ranges, and 3) Sample Collection at Low Velocity Ranges. The overall sampling strategy and soil recommendations that are common to both types of ranges are discussed in the General Sampling Strategy subsection. Recommendations that are specific to the two types of ranges are provided in the second and third subsections.

***General Sampling Strategy.*** Sampling may be required to support several different management strategies at a shooting range. The initial visual inspection of the site and a review of operating records may indicate that sufficient spent ammunition exists, making removal and recycling cost effective. For small ranges, particularly pistol and rifle ranges, it may be cost effective to remove or remediate contaminated material rather than conduct a risk assessment. In this case, an initial sampling maybe conducted to determine the amount and type of sampling data required to characterize the material for treatment or disposal, followed by a screening to ensure the remaining soil does not pose a potential risk. Even when recycling will not be a primary management strategy for the site, it is recommended to reclaim spent ammunition from the range prior to collection of data to support a risk assessment in order to avoid duplication of effort. The risk assessment may be conducted to determine the potential for imminent risk, evaluate the risk posed by residual levels of lead in soil, or to develop further management strategies. A first step in the risk assessment is to define the exposure area(s) for

the site. Exposure areas should be small enough to reflect an area of repeated site use by a hypothetical individual. Additional exposure areas may need to be defined for agricultural activities, including crop planting, maintenance and harvesting, and grazing.

The following recommendations for sampling have been developed to produce data that are adequate for use with the IEUBK and/or ALM. Given the primary exposure pathways (i.e., incidental ingestion of soil), the objectives of the sampling effort should include producing precise estimates of the exposure point concentration (EPC) for lead in soil. Sampling designs for outdoor shooting ranges should be appropriate for estimating the mean of skewed distributions (Cochran, 1977, p.40; Chen, 1995). Data from skeet and trap ranges indicate the distribution of lead is typically positively skewed, with concentrations in sieved samples ranging from <1 to 161,000 ppm and coefficients of skewness ranging from 1.0 to 8.3 (Dragun Corp., 1992; EA, 1994, 1995; E&E, 1997; Murray et al., 1997; TTNUS, 2001). Data from rifle and pistol ranges indicate the range of lead concentration may vary from background levels to 3.9% lead (39,000 ppm), by weight (Dragun Corp., 1992; EEA, 1992).

Sampling depth should be appropriate for the exposure scenario(s) that are to be considered in the risk assessment. Typically, this will dictate that samples be collected from the surficial soils (i.e., 0–1” depth interval) to assess current exposure scenarios. To assess the risk for future exposure scenarios it may be appropriate to also collect samples at depth. If a large number of samples from different depth intervals are planned, it is suggested to evaluate the correlation in concentration/shot density between depth intervals to determine if samples from the different depth intervals can be combined to reduce sampling costs (U.S. EPA, 2000).

Site conditions (e.g., wetlands) may restrict access to some areas of the site or may increase sampling costs. The sampling plan should account for this to avoid introducing bias into the estimate of the EPC. Samples of other media (e.g., surface water) should be collected as appropriate for the exposure scenarios considered in the risk assessment. Assistance of a statistician to develop the sampling design is recommended.

**Sample Preparation.** Sieving of soil samples, to evaluate risks, is recommended for two reasons: the fine particle size fraction (<250  $\mu\text{m}$ ) is the primary source of soil ingestion (U.S. EPA, 2000) and should be used in predicting risk to humans for the incidental soil ingestion pathway; secondly, sieving will remove lead shot and large bullet fragments from the sample, which are not likely to be ingested inadvertently by humans. It is recommended that soil samples be sieved twice, first with a No. 4 (4.75 mm) or No. 10 (2.00 mm) sieve to remove bulk debris, then with a No. 60 (250  $\mu\text{m}$ ) sieve, or smaller sieve size (U.S. EPA, 2000). The No. 4 sieve size recommendation is based on the maximum size shot that is typically used on skeet, trap, and sporting clay ranges (No. 7½), which has a diameter of 2.41 mm. Table B-1 lists the diameters of the pellets for different shot sizes and the size of the openings for different sieve sizes. The portion of the sample that passes through the No. 4 or No. 10 sieve, but retained on the No. 60 sieve, is the ‘coarse fraction’; the portion passing through the No. 60 sieve is the ‘fine fraction’ (U.S. EPA, 2000). The portion of the sample passing through the first sieve (i.e., No. 4 or No. 10) may be referred to as the ‘total’ sample (i.e., coarse + fine fractions). The ‘total’ soil concentration may be appropriate for predicting risks to ecological receptors and for

predicting risks to humans for future exposure scenarios.

It may be possible to reduce sampling costs by developing a relationship between the coarse and fine sample fractions and use the concentration in one fraction to predict the concentration in the other fraction (U.S. EPA, 2000). However, the coarse fraction will contain fragments from bullets and shot, which will tend to dominate the concentrations measured in the sample. Unless the lead fragments and shot are distributed uniformly across the site it is unlikely that the lead concentration between the coarse and fine fractions will be highly correlated. Increasing the volume of the sample or collecting composite samples may help to improve the correlation.

**Analytical Methods.** Samples of the fine fraction should be analyzed for total lead to predict the risk from incidental ingestion of soil. The total fraction should be analyzed for total lead to predict the risks from exposure that may occur after the bullet fragments and shot have undergone additional weathering. Solid waste test method 7421 is recommended for measuring the concentration of lead in soil when using fixed-based laboratories for analysis (EPA, 1986b). The use of field-based devices (e.g., X-ray Fluorescence [XRF]) to measure the concentration of lead may be cost-effective and decrease the time to site cleanup. EPA guidance on the use of field-based methods (U.S. EPA, 2001c) should be consulted prior to developing a sampling plan for the site.

**Sample Collection at High Velocity Ranges.** The horizontal boundaries of active or recently closed pistol and rifle ranges should be fairly obvious; the boundaries of pistol and rifle ranges that have been abandoned for longer periods of time may not be readily apparent from visual inspection alone. Soil samples should be collected from the berm and the rest of the shooting range(s). Samples of other media should be collected, as appropriate, given the exposure scenarios considered in the risk assessment.

Soil sample locations should be determined using random sampling methods that provide adequate coverage of the site, e.g., using systematic or stratified random sampling methods (Gilbert, 1987).

For pistol and rifle ranges, either of these sampling designs may be implemented by sampling on a rectangular or triangular grid. Jacketed bullets typically travel deeper into berms than unjacketed bullets (EA, 1996); the sampling plans for military ranges and other ranges that use jacketed bullets should take this into consideration.

**Sample Collection at Low Velocity Ranges.** Determining the extent of the area potentially affected by skeet and trap ranges are usually more difficult than it is for pistol and rifle ranges. This is particularly true for closed ranges. Whether the range is active or closed, records of site operations, the location of structures on the range (e.g., traphouse) and information gathered from site inspections can help to prepare a preliminary site layout upon which an initial sampling design can be based.

For skeet and trap ranges, a radial grid, with the origin located behind the shooting positions, may produce a more efficient systematic or stratified random sampling design (e.g., EA, 1995). The highest concentrations of lead are typically found in the top 6–8 inches of soil (EA, 1994, 1995; E&E, 1997;

Murray, 1997); however, elevated concentrations of lead have been detected at 24 inches below grade in skeet and trap range shotfall zones (EEA, 1992; EA, 1995; Murray, 1997).

### **Lead Risk Modeling Recommendations**

This section provides recommendations specific to ranges. General guidance on lead risk assessment, lead models and model documentation can be downloaded from the [TRW web site](#) (U.S. EPA, 2002d). The TRW recommends the use of the IEUBK and ALM to predict risk to children and adults, respectively (U.S. EPA, 1994b, 1996). The models are intended to predict risk to humans from exposure to lead that is continuously distributed in various media, including soil. Spent lead ammunition in soil poses a potential risk to humans in two forms: 1) as lead adsorbed or absorbed to soil particles, and as very fine lead particles; and, 2) as lead adsorbed/absorbed to larger soil particles, and as lead shot and bullet fragments. Small soil particles (and by analogy, small lead particles), particularly those <250  $\mu\text{m}$  in size, are the primary source of soil ingestion (EPA, 2000). Lead shot, bullets and large bullet fragments, and large soil particles contaminated with lead, are not likely to be ingested by humans, but represent a source of lead that may be released to the environment through weathering processes. The first form of spent lead ammunition should be considered under the current exposure scenario, while both forms of spent lead ammunition should be considered under future use scenarios.

The IEUBK and the ALM require the user to input an estimate for the EPC for soil; the IEUBK also provides the user with the option of inputting estimates of the EPC for other media: dust, air, drinking water, and diet (e.g., consumption of game and fish with elevated lead concentrations due to exposure to spent lead ammunition). Recommendations are limited here to providing estimates of the EPC for soil; recommendations for other media and model parameters are provided in the IEUBK User's Manual and other guidance that is available on the [TRW website](#) (U.S. EPA, 2002d) and the reference list of this paper.

***Recommendations for Estimating the Exposure Point Concentration Term.*** When interpreting analytical data for a shooting range, it is important to distinguish between high and low velocity ranges. As described in beginning of the Site Characterization and Risk Assessment section, soil on high velocity ranges tend to contain very fine particles of lead in addition to bullets and bullet fragments. Soil on low velocity ranges will tend to contain whole and partially decomposed lead pellets. The presence of one bullet, bullet fragment, or lead shot in a soil sample will result in very high measurement of lead concentration which may not yield an accurate prediction of risk for current exposure scenarios (but may be appropriate for future exposure scenarios).

The use of geostatistics may be useful for shooting ranges, particularly skeet and trap ranges, where soil concentrations often exhibit spatial patterns. Geostatistical estimators are capable of exploiting these spatial patterns (i.e., spatial autocorrelation) to determine the extent of contaminated soil, and to produce more precise estimates of the EPC. Another advantage of geostatistical estimators is they can be used with data that have been collected by random and/or non-random sampling methods. Finally, geostatistics can take advantage of the correlation between different types of measurements (e.g., soil concentration and shot count) to obtain more precise estimates of the EPC.

***Recommendations for Adjusting Bioavailability.*** The TRW does not recommend changing the default value for bioavailability without the collection and TRW review of good site-specific data to support such a change. Bioavailability has been shown to be related to lead speciation and soil particle size (U.S. EPA, 1999). While site specific data on lead speciation (e.g., from decomposition of spent ammunition) and particle size are not considered by EPA to be an adequate basis for adjusting the bioavailability variable in the IEUBK or ALM (U.S. EPA, 1999), this information can be used to decide if *in vivo* bioassays are likely to produce estimates of bioavailability that differ substantially from the IEUBK default value. Guidance on the bioavailability variable is available from the [TRW website](#) (U.S. EPA, 1999). Based on the available literature on lead speciation in soil on shooting ranges, it appears the default values used in the IEUBK and ALM are appropriate for assessing risks from exposure to soils located on shooting ranges.

***Recommendations for Exposure Scenarios.*** Land use adjacent and near shooting ranges should be considered when developing current and future exposure scenarios. For currently operating sites, the populations of primary concern are residents of adjacent residential properties, residents and farm workers on adjacent agricultural properties, and workers who are employed on adjacent commercial properties. For future use scenarios, the populations of primary concern depend upon the proposed site usage.

#### **BEST MANAGEMENT PRACTICES AND RECYCLING**

Guidance for implementing best management practices on small arms outdoor shooting ranges is available from the EPA ([EPA, 2001a](#)). Implementing best management practices (BMPs) on ranges decreases the risk of spent ammunition to the environment by recycling spent ammunition; containing lead shot, bullets, and fragments; preventing migration of lead to surface water and groundwater; and keeping records of site operations (U.S. EPA, 2001a). Site conditions (e.g., wetlands, mud, steep slopes, wooded areas) will affect the feasibility of removing spent ammunition from ranges.

## References

- ATSDR. 1999. *Toxicological Profile for Lead (Update)*. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, Agency for Toxic Substances and Disease Registry. Atlanta, GA. July. Available online at: <http://www.atsdr.cdc.gov/toxprofiles/tp13.html>.
- Baer, K.N., D.G. Hutton, R.L. Boer, T.J. Ward, and R.G. Stahl. 1995. Toxicity Evaluation of Trap and Skeet Shooting Targets to Aquatic Test Species. *Ecotoxicology*. 4:385-92.
- Baldwin, D. 1994. How Far Will a Shotgun Shoot? Gun Club Advisor. Spring. 1994. Available on-line at: [http://www.rangeinfo.org/resource\\_library/facility\\_mngmnt/design/how\\_far\\_will.htm](http://www.rangeinfo.org/resource_library/facility_mngmnt/design/how_far_will.htm).
- Battelle. 1997. Implementation Guidance Handbook: Physical Separation and Acid Leaching to Process Small-Arms Range Soils. Prepared for Naval Facilities Engineering Service Center and U.S. Army Environmental Center. September.
- Bundy, K.J., M. Bricka, and A. Morales. 1996. *An Electrochemical Approach for Investigating Corrosion of Small Arms Munitions in Firing Ranges*. Proceedings of the HSRC/WERC Conference on the Environment. Albuquerque, NM. May 21-23.
- Capital Trap Club. 2001a. American Trapshooting. Ottawa, Canada. April, 4. Available on-line at: <http://www.firearmstraining.ca/capital/americantrap.html>.
- Capital Trap Club. 2001b. American Skeet. Ottawa, Canada. April, 4. Available on-line at: <http://www.firearmstraining.ca/capital/skeet.html>.
- CDC. 1991. *Preventing Lead Poisoning in Young Children*. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention. Atlanta, GA. Available on-line at: <http://wonder.cdc.gov/wonder/prevguid/p0000029/p0000029.asp>.
- Chen, L. 1995. Testing the Mean of Skewed Distributions. *J. Amer. Stat. Assoc.* 90(430):767-772.
- Cochran, W.G. 1977. *Sampling Techniques*. 3<sup>rd</sup> Ed. John Wiley & Sons, New York, NY.
- Dragun Corp. 1992. Phase II Environmental Assessment Report Parcels 32/36 Carleton Sportsmen's Club Sumpter Township, Michigan. October.
- EA Engineering, Science and Technology (EA). 1994. Phase I Site Assessment Southern Lakes Trap and Skeet Club. October 28, 1994.
- EA Engineering, Science and Technology (EA). 1995. Phase II Site Investigation Camp Buckner Skeet and Trap Range. Final. U.S. Military Academy. West Point, NY. August.



EA Engineering, Science and Technology (EA). 1996. *Lead Mobility at Shooting Ranges*. Sporting Arms and Ammunition Manufacturers' Institute, Inc. Newton, CT.

E&E (Ecology and Environment, Inc.). 1997. Preliminary Assessment/Site Inspection for Nahant Marsh Site, Davenport, Iowa. Final. September.

EEA, Inc. 1992. Environmental Assessment and Field Investigation at the Blue Mountain Sportsmen's Center, Town of Cortlandt, Westchester County, New York. Preliminary Draft. November.

Gilbert, R.O. 1987. *Statistical Methods for Environmental Pollution Monitoring*. Van Nostrand Reinhold, New York, NY.

Hall, T., Jr. 2002. Personal communication. July 3.

Henningsen, G., C. Weis, S. Casteel, L. Brown, E. Hoffman, W. Brattin, J. Drexler, and S. Christensen. 1998. Differential Bioavailability of Lead Mixtures from Twenty Different Soil Matrices at Superfund Mining Sites. Abstract. *Toxicol. Sci.* 42(1-s).

Hoffman, D.J., G.H. Heinz, L. Sileo, D.J. Audet, J.K. Campbell, and L.J. LeCaptain. 2000. Developmental Toxicity of Lead-Contaminated Sediment to Mallard Ducklings. *Arch. Environ. Contam. Toxicol.* 39:221-232.

Jorgensen, S. and M. Willems. 1987. The Fate of Lead in Soils: The Transformation of Lead Pellets in Shooting-Range Soils. *Ambio.* 16(1):11-15.

Lin, Z., B. Comet, U. Qvarfort, and R. Herbert. 1995. The Chemical and Mineralogical Behaviour of Pb in Shooting Range Soils from Central Sweden. *Environ. Poll.* 89(3):303-09.

Ma, Wei-chun. 1989. Effect of Soil Pollution with Metallic Lead Pellets on Lead Bioaccumulation and Organ/Body Weight Alterations in Small Mammals. *Arch. Environ. Contam. Toxicol.* 18:617-622.

Major, M. 2003. Personal communication. March 3.

Manninen, S. and N. Tanskanen. 1993. Transfer of Lead from Shotgun Pellets to Humus and Three Plant Species in a Finnish Shooting Range. *Arch. Environ. Contam. Toxicol.* 24:410-414.

Murray, K., A. Bazzi, C. Carter, A. Ehlert, A. Harris, M. Kopec, J. Richardson, and H. Sokol. 1997. Distribution and Mobility of Lead in Soils at an Outdoor Shooting Range. *J. Soil Contam.* 6(1):79-93.

NRA (National Rifle Association). 1999. The Range Source Book. National Rifle Association, Range Department, Field Operations Division. Fairfax, VA. November.

Peddicord, R.K. and J.S. LaKind. 2000. Ecological and Human Health Risks at an Outdoor Firing

Range. *Env. Tox. Chem.* 19(10):2602-13.

Roscoe, D.E., L. Widjeskog, and W. Stansley. 1989. Lead Poisoning of Northern Pintail Ducks Feeding in a Tidal Meadow Contaminated with Shot from a Trap and Skeet Range. *Bull. Environ. Contam. Toxicol.* 42:226-233.

TTNUS (Tetra Tech NUS, Inc.). 2001. Ecologically-Based Remediation Goals for Lead and PAHs in Soil, Site 15, Blue 10 Ordinance Disposal Area, Naval Air Station Cecil Field, Jacksonville, Florida.

U.S. EPA. 1986a. *Air Quality Criteria for Lead*. Vol I. Draft Final. EPA-600/8-83/028aF. June.

U.S. EPA. 1986b. *SW-846: Test Methods for Evaluating Solid Wastes Physical/Chemical Methods, Method 7421, Lead ( Atomic Adsorption, Furnace Technique)*. Revision 0. September. Available on-line at: <http://www.epa.gov/epaoswer/hazwaste/test/pdfs/7421.pdf>.

U.S. EPA. 1994a. *OSWER Directive: Revised Interim Soil Lead (Pb) Guidance for CERCLA Sites and RCRA Corrective Action Facilities*. EPA/540/F-94/043 PB 94-963282. August. Available on-line at: <http://www.epa.gov/superfund/programs/lead/products/oswerdir.pdf>.

U.S. EPA. 1994b. *Guidance Manual for the Integrated Exposure Uptake Biokinetic Model for Lead in Children*. Office of Emergency and Remedial Response. EPA/540/R-93/081 PB 93-9635100. February. Available on-line at: <http://www.epa.gov/superfund/programs/lead/products.htm>.

U.S. EPA. 1996. *Recommendations of the Technical Review Workgroup for Lead for an Interim Approach to Assessing Risks Associated with Adult Exposures to Lead in Soil*. December. Available on-line at: <http://www.epa.gov/superfund/programs/lead/products.htm>.

U.S. EPA, 1999. *IEUBK Model Bioavailability Variable*. EPA-540-F-00-006. OSWER 9285.7-32. October. Available on-line at: <http://www.epa.gov/superfund/programs/lead/products/sspbbioc.pdf>.

U.S. EPA, 2000. *TRW Recommendations for Sampling and Analysis of Soil at Lead (Pb) Sites*. EPA-540-F-00-010. OSWER 9285.7-38. April. Available on-line at: <http://www.epa.gov/superfund/programs/lead/products/sssiev.pdf>.

U.S. EPA, 2001a. *Best Management Practices for Lead at Outdoor Shooting Ranges*. EPA-902-B-01-001. Region 2. January. Available on-line at: <http://www.epa.gov/region2/waste/leadshot/>.

U.S. EPA, 2001b. OSW Memorandum to RCRA Senior Policy Advisors. Subject: *National Guidance on Best Management Practices for Lead at Outdoor Shooting Ranges*. October 10.

U.S. EPA. 2001c. *Integrating Dynamic Field Activities Into the Superfund Response Process: A Guide for Project Managers*. Draft Final. December.

U.S. EPA. 2002a. ECOTOX. Office of Research and Development. U.S. Environmental Protection Agency. Washington, DC. Available on-line at: [http://www.epa.gov/ecotox/ecotox\\_home.htm](http://www.epa.gov/ecotox/ecotox_home.htm).

U.S. EPA. 2002b. *Lead and Human Health*. Technical Review Workgroup. Available on-line at: <http://www.epa.gov/superfund/programs/lead/leadandhumanhealth.htm>.

U.S. EPA. 2002c. Lead Workgroup. Products. Region 2. U.S. Environmental Protection Agency. Washington, DC. Available on-line at: <http://www.epa.gov/superfund/programs/lead/products.htm>.

U.S. EPA. 2002d. Technical Review Workgroup for Lead (TRW) Home Page. U.S. Environmental Protection Agency. Washington, DC. Available on-line at: <http://www.epa.gov/superfund/programs/lead/index.htm>.

U.S. FWS. 1999. *Migratory Bird Hunting*. 50 CFR 20.21(j).

U.S. FWS. 2001. *Refuge-Specific Regulations for Fishing and Hunting*. 50 CFR 32.2(k).

Vargas, C. 1996. Design Criteria for Shooting Ranges. Third National Shooting Range Symposium. Available on-line at: [http://www.rangeinfo.org/resource\\_library/facility\\_mngmnt/safety/design\\_criteria.htm](http://www.rangeinfo.org/resource_library/facility_mngmnt/safety/design_criteria.htm).

Vyas, N.B., J.W. Spann, G.H. Heinz, W.N. Beyer, J.A. Jaquette, and J.M. Mengelkoch. 2000. Lead Poisoning of Passerines at a Trap and Skeet Range. *Environ. Poll.* 107:159-166.

**Table A-1. On-line Sources of Information Related to Outdoor Small Arms Shooting Ranges**

Agency/Organization	Main Web Address	Topic	Web Address
Fish and Wildlife Service	<a href="http://www.fws.gov/">http://www.fws.gov/</a>	Hunting information	<a href="http://birds.fws.gov/Laws.htm">http://birds.fws.gov/Laws.htm</a>
		Fish and Wildlife Management Offices - State, Territorial, and Tribal	<a href="http://offices.fws.gov/statelinks.html">http://offices.fws.gov/statelinks.html</a>
Environmental Protection Agency	<a href="http://www.epa.gov/">http://www.epa.gov/</a>	Envirosense PRO-ACT Factsheet on Lead Contamination In Soil at Small Arms Firing Ranges	<a href="http://es.epa.gov/program/p2dept/defense/airforce/2818.html">http://es.epa.gov/program/p2dept/defense/airforce/2818.html</a>
Environmental Protection Agency	<a href="http://www.epa.gov/">http://www.epa.gov/</a>	Military Munitions Rule, Federal Register: February 12, 1997. Volume 62, Number 29. Page 6621-6657	<a href="http://www.epa.gov/docs/fedrgstr/EPA-WASTE/1997/February/Day-12/f3218.htm">http://www.epa.gov/docs/fedrgstr/EPA-WASTE/1997/February/Day-12/f3218.htm</a>
Environmental Protection Agency (EPA) - Region 2	<a href="http://www.epa.gov/">http://www.epa.gov/</a>	Best Management Practices for Lead at Outdoor Shooting Ranges	<a href="http://www.epa.gov/region2/waste/leadshot/">http://www.epa.gov/region2/waste/leadshot/</a>
PROACT (Department of Defense)	<a href="http://www.afcee.brooks.af.mil/pro-act/PRO-ACThome.asp">http://www.afcee.brooks.af.mil/pro-act/PRO-ACThome.asp</a>	Factsheet on lead contamination in soil at small arms firing ranges	<a href="http://www.afcee.brooks.af.mil/pro-act/fact/june98a.asp">http://www.afcee.brooks.af.mil/pro-act/fact/june98a.asp</a>

<b>Agency</b>	<b>Main Web Address</b>	<b>Topic</b>	<b>Web Address</b>
Alabama Department of Conservation and Natural Resources	<a href="http://www.dcnr.state.al.us/agfd/index.html">http://www.dcnr.state.al.us/agfd/index.html</a>	Legal arms and ammunition for hunting	<a href="http://www.dcnr.state.al.us/agfd/arms.html">http://www.dcnr.state.al.us/agfd/arms.html</a>
Florida Fish and Wildlife Conservation Commission	<a href="http://floridaconservation.org/">http://floridaconservation.org/</a>	Public shooting ranges	<a href="http://floridaconservation.org/huntered/ranges.html">http://floridaconservation.org/huntered/ranges.html</a>
		Hunting information and regulations	<a href="http://www.wld.fwc.state.fl.us/hunting/default.html">http://www.wld.fwc.state.fl.us/hunting/default.html</a>
Idaho Fish and Game	<a href="http://www2.state.id.us/fishgame/common/50list.htm">http://www2.state.id.us/fishgame/common/50list.htm</a>	50 State Fish and Game Agencies List	<a href="http://www2.state.id.us/fishgame/common/50list.htm">http://www2.state.id.us/fishgame/common/50list.htm</a>
Massachusetts Division of Fisheries, Wildlife and Environmental Law Enforcement	<a href="http://www.state.ma.us/dfwele/dpt_toc.htm">http://www.state.ma.us/dfwele/dpt_toc.htm</a>	Lead shot in the environment	<a href="http://www.state.ma.us/dep/files/pbshot/pb_shot.htm">http://www.state.ma.us/dep/files/pbshot/pb_shot.htm</a>
		Wildlife recreation information; hunting and fishing laws	<a href="http://www.state.ma.us/dfwele/dfw/dfwrec.htm#LAWS">http://www.state.ma.us/dfwele/dfw/dfwrec.htm#LAWS</a>
Michigan Department of Environmental Quality	<a href="http://www.michigan.gov/deq/">http://www.michigan.gov/deq/</a>	Environmental regulations affecting shooting ranges	<a href="http://www.michigan.gov/deq/1,1607,7-135-3585_4127_13090-25492--,00.html">http://www.michigan.gov/deq/1,1607,7-135-3585_4127_13090-25492--,00.html</a>
Ohio Environmental Protection Agency	<a href="http://www.epa.state.oh.us/">http://www.epa.state.oh.us/</a>	List of lead reclaimers	<a href="http://www.epa.state.oh.us/dhwm/leadrecy.htm">http://www.epa.state.oh.us/dhwm/leadrecy.htm</a>
Tennessee Wildlife Resources Agency	<a href="http://www.state.tn.us/twra/index.html">http://www.state.tn.us/twra/index.html</a>	Legal hunting equipment and methods	<a href="http://www.state.tn.us/twra/hunt001a3.html">http://www.state.tn.us/twra/hunt001a3.html</a>
National Shooting Sports Foundation (NSSF)	<a href="http://www.huntinfo.org/">http://www.huntinfo.org/</a>	Summaries of every state's hunting opportunities and regulations and links to state fish and game websites	<a href="http://www.huntinfo.org/">http://www.huntinfo.org/</a>

<b>Agency</b>	<b>Main Web Address</b>	<b>Topic</b>	<b>Web Address</b>
		Search page to find shooting ranges	<a href="http://www.wheretoshoot.org/">http://www.wheretoshoot.org/</a>
National Association of Shooting Ranges (NASR)	<a href="http://www.rangeinfo.org/">http://www.rangeinfo.org/</a>	Includes references and sources of information on outdoor shooting ranges	<a href="http://www.rangeinfo.org/">http://www.rangeinfo.org/</a>
National Sporting Clays Association	<a href="http://WWW.NSSA-NSCA.COM/nsca/index.htm">http://WWW.NSSA-NSCA.COM/nsca/index.htm</a>	Search page to find shooting ranges	<a href="http://WWW.NSSA-NSCA.COM/nsca/index.htm">http://WWW.NSSA-NSCA.COM/nsca/index.htm</a>
		Links to shooting-related sites	<a href="http://www.nssa-nsca.com/common/shooting_sites.htm">http://www.nssa-nsca.com/common/shooting_sites.htm</a>
Sporting Arms and Ammunition Manufacturer's Institute, Inc. (SAAMI)	<a href="http://www.saami.org/">http://www.saami.org/</a>	General source of information on shooting ranges	<a href="http://www.saami.org/">http://www.saami.org/</a>
Miscellaneous	<a href="http://dir.yahoo.com/Recreation/Outdoors/Hunting/Organizations/">http://dir.yahoo.com/Recreation/Outdoors/Hunting/Organizations/</a>	List of hunting organizations	<a href="http://dir.yahoo.com/Recreation/Outdoors/Hunting/Organizations/">http://dir.yahoo.com/Recreation/Outdoors/Hunting/Organizations/</a>
National Wild Turkey Federation	<a href="http://www.shooting-hunting.com/index.html">http://www.shooting-hunting.com/index.html</a>	search page for shooting ranges	<a href="http://www.shooting-hunting.com/results.html?Keywords=Shooting+Range">http://www.shooting-hunting.com/results.html?Keywords=Shooting+Range</a>

**Table B-1. Shot Size and EPA-Recommended Sieve Sizes  
for Use at Small Arms Outdoor Firing Ranges**

Shot Size	Pellet Diameter	
	Inches	Millimeters
Buckshot		
No. 000 - No. 2	.36 - .27	9.14 - 6.86
No. 3	.25	6.35
No. 4	.24	6.10
Shot		
F	.22	5.59
T	.20	5.08
BBB	.19	4.83
<b>A NO. 4 SIEVE (0.19 INCH/4.76 MM OPENINGS) WILL REMOVE THE SHOT SIZES LISTED ABOVE.</b>		
BB	.18	4.57
1	.16	4.06
2	.15	3.81
3	.14	3.56
4	.13	3.30
5	.12	3.05
6	.11	2.79
7	.10	2.54
7 ½	.095	2.41
8	.09	2.29
8 ½	.085	2.16
9	.08	2.03
<b>A NO. 10 SIEVE (0.08 INCH/2.00 MM OPENINGS) WILL REMOVE THE SHOT SIZES LISTED ABOVE</b>		
<b>A NO. 60 (0.25 MM OPENINGS) SIEVE, OR SMALLER SIEVE SIZE, IS RECOMMENDED TO PREPARE THE 'FINE' SOIL FRACTION FOR ANALYSIS (SEE 'SAMPLE PREPARATION' SECTION)</b>		
<p>Note: Shot size is generally limited to a maximum of no. 7 ½ for trap and sporting clay use, and a maximum of no. 7 ½ and minimum of no. 9 for skeet shooting.</p>		